



Orthopaedic Associates of Zanesville
Sports Medicine Division Physical Therapy Protocols
Brad E. Brautigan MD

Rehabilitation Guidelines for ACL Reconstruction with Meniscal Repair (All Inside)

The intent of this protocol is to provide the clinician with a general guideline of the post-operative course of the patient that has undergone the above stated procedure. It is by no means intended to substitute for one's clinical decision making regarding the progression of a patient's post-operative course based on their physical exam findings, individual progress, and/or the presence of possible post-operative complications. If a clinician requires assistance in the progression, please contact the referring surgeon's office.

0-2 Weeks:

- TTWB, Brace locked at 0 degrees for ambulation and sleeping, unlocked for rehab
- ROM: 0-90° with emphasis on full extension
- Patella mobilization
- SLR supine with brace locked at 0 degrees, Quad Sets, VMO adduction
- Ankle Pumps, Heel slides
- Short crank (90mm) ergometry

2-6 Weeks:

- Begin WBAT. Unlock Brace for Weight Bearing
- No weight bearing past 90° for ACL with meniscal repair
- D/C crutches when gait is non-antalgic (two weeks with meniscal repair)
- ROM: 0-125 degrees (Maintain full extension)
- Active knee extension to 40 degrees
- Standard (170mm) ergometry (when knee ROM > 115 degrees)
- Leg Press (80-0 degree arc)
- Mini Squats / Weight Shifts
- Proprioception training
- Initiate Step Up program
- Avoid Tibial Rotation until 6 weeks

6-14 Weeks:

- D/C Brace and wean from crutches
- Progressive Squat program
- Initiate Step Down program
- Leg Press, Lunges
- Isotonic Knee Extensions (90-40 degrees, closed chain preferred)
- Agility exercises (sport cord)
- Versaclimber/Nordic Track
- Retrograde treadmill ambulation

14-22 weeks:

- Begin forward running (treadmill) program when 8" step down satisfactory
- Continue Strengthening & Flexibility program
- Advance Sports-Specific Agility Drills
- Start Plyometric program

>22 weeks:

- Advance Plyometric program, Return to Sport (MD Directed)
- May require Functional Sports Assessment (FSA) 5-6 months post op for clearance to return to sport