

Benjamin R. Pulley, MD
*Orthopaedic Trauma Surgeon
& Joint Reconstruction Specialist*
Orthopaedic Associates of Zanesville
2854 Bell Street
Zanesville, OH 43701
740-454-3273

Postoperative Instructions after SURGICAL FIXATION OF TIBIAL PLATEAU FRACTURE

1. Activity instructions:
 - a. No weight bearing with your surgical lower extremity.
 - b. You must use crutches or a walker to keep weight off of your surgical lower extremity when moving around.
 - c. If you were provided with a hinged knee brace, wear it at all times (other than showering).
 - d. You may work on gentle range of motion of your knee while wearing the hinged knee brace.
2. Surgical dressing instructions:
 - a. You may remove the Ace wrap and cotton dressing on postoperative day #3.
 - b. You have a waterproof dressing beneath the Ace wrap and cotton. The waterproof dressing may remain in place until your first postoperative follow-up appointment in the office.
 - c. Once you have removed the Ace wrap and cotton dressing, you may shower when desired, but do not submerge the surgical lower extremity in a bath tub.
3. Prevention of deep venous thrombosis (blood clot):
 - a. If you **do not usually take any other blood thinning medications**, then you will start taking **Aspirin 81 mg by mouth twice daily** on the morning of postoperative day #1 and continue this for 4 weeks.
 - b. If you **already take a blood thinning medication** at baseline, then your instructions are as follows: _____

4. Pain control:
 - a. The best way to control pain is to control swelling. The best way to control swelling is to elevate the surgical lower extremity above the level of your heart and apply ice often.
 - b. An electronic prescription has been sent to your pharmacy for a 1 week supply of narcotic pain medication. Narcotic pain medication is sedating and constipating. Increase your fluid intake and use over-the-counter stool softeners as needed to help with constipation. Use the narcotic pain medication when your pain is most severe. Wean yourself off of the narcotic pain medication as soon as possible.

- c. You may take acetaminophen in addition to, or in substitution for your narcotic pain medication, but do not exceed 3000 mg of acetaminophen in a 24 hour period from all sources.
 - d. You may take non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen in addition to, or in substitution for your narcotic pain medication. Some patients have been advised by their family physician to avoid NSAIDs. If you are unsure, call your family physician.
 - e. It is safe to alternate between narcotic pain medication, acetaminophen, and NSAIDs.
5. Reasons to call the office:
- a. Foul odor or excessive drainage from your surgical dressing.
 - b. Increasing pain not relieved by pain medication.
 - c. New numbness or tingling.
 - d. Signs or symptoms of systemic illness (fever, chills, nausea, generalized weakness or body aches, etc.).
6. Follow-up:
- a. A 2 week follow-up appointment was made in the office on the day that you scheduled your surgery. If you do not know the date of the follow-up appointment, please call the office at your convenience for a reminder.